



Crops(s) Avocado and Mango

Disease(s) Anthracnose (*Colletotrichum* spp.)

Resistance management strategy for:

Group 3*	Demethylation inhibitors (DMI);
Group 7+11	Succinate dehydrogenase inhibitors (SDHI) + Quinone
	outside inhibitor (QoI); and
Group 11	Quinone outside inhibitor (QoI)

Guidelines:

- If applying Group 11 (including 7+11) fungicides, do not apply more than two consecutive sprays before changing to another group. Do not apply more than three Group 11 sprays per season within the field. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another Group 11.
- 2. ***Do not** apply more than four **Group 3** sprays in a season. Apply no more than two consecutive sprays of a **Group 3** fungicide alone.
- 3. No specific resistance management strategy has been developed for low-risk fungicides, including those in **Group M** and **BM**. These products should be included in a management strategy as per label recommendations.

* Appropriate in mangoes only

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 30 June 2022. All previous versions of this strategy are now invalid.

